

## The Cytotoxic Activity of *Annona muricata* Leaf Oil from Badagary, Nigeria

Moses S. Owolabi, Akintayo Lanre Ogundajo, Noura S. Dosoky, and William N. Setzer

### ABSTRACT

The leaf essential oil of *Annona muricata*, collected from Badagary, Nigeria, was obtained by hydrodistillation and analyzed by gas chromatography – mass spectrometry. The leaf oil was dominated by (*E*)-caryophyllene (38.9%) and eugenol (30.2%), with lesser amounts of  $\alpha$ -humulene (4.3%),  $\delta$ -cadinene (6.0%), and caryophyllene oxide (5.0%). *A. muricata* leaf oil showed notable *in-vitro* cytotoxic activity on MCF-7 cells (99.2% kill at 100  $\mu$ g/mL), which can be attributed to the major components.

**Keywords:** *Annona muricata*, Essential Oil Composition, Cytotoxicity, Caryophyllene, Eugenol.

AJEONP 2013; 1 (1): 1-3  
© 2013 AkiNik Publications  
Received 17-7-2013  
Accepted: 28-7-2013

**Moses S. Owolabi**  
Department of Chemistry, Lagos State  
University P.M.B. 001, Ojo, Lagos,  
Nigeria.

**Akintayo Lanre Ogundajo**  
Department of Chemistry, Lagos State  
University P.M.B. 001, Ojo, Lagos,  
Nigeria.

**Noura S. Dosoky**  
Department of Chemistry, University  
of Alabama in Huntsville Huntsville,  
Alabama 35899,  
USA

**William N. Setzer**  
Department of Chemistry, University  
of Alabama in Huntsville Huntsville,  
Alabama 35899,  
USA

**Correspondence:**  
**Moses S. Owolabi**  
Department of Chemistry, Lagos  
State University P.M.B. 001, Ojo,  
Lagos, Nigeria.  
E-Mail: [sunnyconcept2007@yahoo.com](mailto:sunnyconcept2007@yahoo.com)

### 1. Introduction

*Annona* is a genus of the Annonaceae, of which there are about 129 species distributed mainly in Tropical and subtropical region including part of the Caribbean, Central and South America, Africa, Asia and Australia <sup>[1]</sup>. Seven species and one hybrid are grown for domestic/commercial use <sup>[2]</sup>. *Annona muricata* L., commonly known in English speaking countries as ‘soursop’ and ‘ebo’ in Yoruba, is an upright, low-branching tree reaching 8 to 10 meters <sup>[3-5]</sup>. The tree has green, glossy evergreen leaves, and the flowers appear anywhere on the trunk or any branch <sup>[6]</sup>. Traditionally, the leaves are used for headaches, insomnia, cystitis, liver problems, diabetes, hypertension and as an anti-inflammatory, antispasmodic and antidiarrheal <sup>[7,8]</sup>. In this work, we present the chemical composition and cytotoxic activity of the leaf essential oil of *A. muricata* growing in Badagary, Nigeria.

### 2. Materials and Methods

Fresh leaves of *Annona muricata* were collected in October, 2011, from Badagary, Lagos state, Nigeria. The plant was taxonomically identified and authenticated at the Herbarium of the Department of Botany of the University of Lagos. Prior to hydrodistillation, the plant was air-dried for three days and pulverized. A sample (350 g) of *A. muricata* was subjected to hydrodistillation in a Clevenger-type apparatus (British Pharmacopoeia 1980) for 4 h. The yield of oil was 0.73% on a pulverized weight basis. The oil was dried over anhydrous sodium sulfate and stored in a sealed vial under refrigeration prior to analysis. The leaf essential oil of *A. muricata* was analyzed by GC-MS as previously described <sup>[9]</sup>. The leaf oil composition is summarized in Table 1. *In-vitro* cytotoxicity evaluation of *A. muricata* oil and essential oil components against MCF-7 human breast adenocarcinoma cells was carried out using the MTT assay as previously described <sup>[10]</sup>.

### 3. Results and Discussion

GC-MS analysis of the leaf essential oil of *A. muricata* (Table 1) revealed 19 identifiable components comprising 100% of the composition. The oil was composed largely of the sesquiterpene hydrocarbons (*E*)-caryophyllene (38.9%),  $\delta$ -cadinene (6.0%), and  $\alpha$ -humulene (4.3%), as well as the phenylpropanoid eugenol (30.2%) and the sesquiterpenoid caryophyllene oxide (5.0%).

*A. muricata* from Brazil was somewhat different, dominated by spathulenol (14.3%), bicyclogermacrene (9.8%), and linalool (8.7%)<sup>[14]</sup>.

The Nigerian *A. muricata* leaf oil showed notable *in-vitro* cytotoxic activity against MCF-7 human adenocarcinoma cells with 99.2 ± 0.1% killing at 100 µg/mL. The cytotoxic activity can be attributed to the major components in the oil. (*E*)-Caryophyllene<sup>[15-18]</sup>, eugenol<sup>[18,19]</sup>, δ-cadinene<sup>[15]</sup>, caryophyllene oxide<sup>[17,18,20]</sup>, α-humulene<sup>[17,18,21,22]</sup>, and β-elemene<sup>[23-25]</sup> have shown cytotoxic activity against several different cell lines. Against MCF-7 cells, (*E*)-caryophyllene, eugenol, caryophyllene oxide, and α-humulene

had *IC*<sub>50</sub> values of 38.8 ± 5.4, 97.1 ± 11.7, 23.0 ± 1.3, and 22.1 ± 1.7 µg/mL, respectively.

#### 4. Conclusions

*A. muricata* leaf oil from Nigeria was rich in (*E*)-caryophyllene and eugenol; the high eugenol content differentiating it from oils originating from other geographical locations. The leaf oil was cytotoxic, which is attributed to the cytotoxicities of each of the major components.

**Table 1:** Chemical composition of *Annona muricata* leaf essential oil from Nigeria.

RI	Compound	%	RI	Compound	%
941	α-Pinene	0.8	1453	α-Humulene	4.3
978	β-Pinene	0.3	1481	γ-Muurolene	0.8
1024	p-Cymene	0.6	1497	Bicyclogermacrene	2.0
1058	γ-Terpinene	0.4	1501	α-Muurolene	2.1
1176	Terpinen-4-ol	0.2	1505	Germacrene A	3.0
1190	α-Terpineol	0.4	1524	δ-Cadinene	6.0
1356	Eugenol	30.2	1583	Caryophyllene oxide	5.0
1375	α-Copaene	0.2	1641	τ-Muurolol	0.7
1392	β-Elemene	3.5	1655	α-Cadinol	0.7
1419	( <i>E</i> )-Caryophyllene	38.9		Total Identified	100.0

#### 5. Reference:

- Mabberley DJ. Mabberley's Plant Book, 3<sup>rd</sup> Ed. Cambridge University Press, Cambridge, UK, 2008, 49.
- International Centre for Underutilized Crops. Fruits for the future. Newsletter 5. March. Southampton, UK: Institute of Irrigation and Development Studies, University of Southampton, 2002. <http://www.icuc-iwmi.org/files/News/Resources/Factsheets/annona.pdf>. 15 August, 2008.
- Popenoe W. Manual of Tropical and Subtropical Fruits. Hafner Press, New York, 1920, 8.
- Mowry H, Toy LR Wolfe HS. Miscellaneous tropical and subtropical Florida fruits. Florida Agricultural Experiment Station, Bulletin 156, 1953.
- Morton JF. Fruits of Warm Climates. Florida Flair Books, Boynton Beach, Florida, 1987, 75-80.
- Morton JF. The soursop, or guanabana (*Annona muricata* Linn.). Proceedings of the Florida State Horticultural Society, 1966; 79: 355-366.
- Di Stasi LC, Hiruma-Lima CA. Plantas Mediciniais na Amazônia e na Mata Atlântica, 2<sup>nd</sup> Ed. Editora UNESP, São Paulo, Brazil, 2002, 87-112.
- Sousa MP, Matos MEO, Matos FJA, Machados MIL, Craveiro AA. Constituintes Químicos Ativos e Propriedades Biológicas de Plantas Mediciniais Brasileiras, 2<sup>nd</sup> Ed. Editora UFC, Fortaleza, Brazil, 2004, 281-283.
- Owolabi MS, Moronkola A, Lajide L, Palazzo MC, Setzer WN. Chemical composition of the seed volatiles of *Aframomum sceptrum* (Oliv. & T. Hanb.) K. Schum. from Nigeria. Journal of Essential Oil-Bearing Plants, 2010; 13: 753-758.
- Moriarity DM, Bansal A, Cole RA, Takaku S, Haber WA, Setzer WN. Selective cytotoxic activities of leaf essential oils from Monteverde, Costa Rica. Natural Product Communications, 2007; 2: 1263-1268.
- Pélissier Y, Marion C, Kone D, Lamaty G, Menut C, Bessière JM. Volatile components of *Annona muricata* L. Journal of Essential Oil Research, 1994; 6: 411-414.
- Boyom FF, Zollo PHA, Menut C, Lamaty G, Bessière, JM. Aromatic plants of tropical central Africa. Part XXVII. Comparative study of volatile constituents of five Annonaceae species growing in Cameroon. Flavour and Fragrance Journal, 1996; 11: 333-338.
- Kossouh C, Moudachirou M, Adjakidje V, Chalchat JC, Figueredo G. Essential oil chemical composition of *Annona muricata* L. leaves from Benin. Journal of Essential Oil Research, 2007; 19: 307-309.
- Moreira RRD, Simeí NA, Santos LE, Cavaleiro C. Gas chromatography (GC) and gas chromatography-mass spectroscopy (GC/MS) analysis of the essential oil from leaves of *Annona muricata* L. XVII Simpósio de Plantas Mediciniais do Brasil, Cuiabá, Brazil, 2002.
- Kubo I, Morimitsu Y. Cytotoxicity of green tea flavor compounds against two solid tumor cells. Journal of Agricultural and Food Chemistry, 1995; 43: 1626-1628.
- Prashar A, Locke IC, Evans CS. Cytotoxicity of clove (*Syzygium aromaticum*) oil and its major components to human skin cells. Cell Proliferation, 2006; 39: 241-248.
- Schmidt JM, Noletto JA, Vogler B, Setzer WN. Abaco bush medicine: Chemical composition of the essential oils of four aromatic medicinal plants from Abaco Island, Bahamas. Journal of Herbs, Spices and Medicinal Plants, 2006; 12(3): 45-65.
- Wright BS, Bansal A, Moriarity DM, Takaku S, Setzer WN. Cytotoxic leaf essential oils from Neotropical Lauraceae: Synergistic effects of essential oil components. Natural Product Communications, 2007; 2: 1241-1244.
- Atsumi T, Tonosaki K, Fujisawa S. Induction of early apoptosis and ROS-generation activity in human gingival fibroblasts (HGF) and human submandibular gland carcinoma (HSG) cells treated with curcumin. Archives of Oral Biology, 2006; 51: 913-921.
- Jun NJ, Mosaddik A, Moon JY, Jang KC, Lee DS, Ahn KS, Cho SK.

- Cytotoxic activity of  $\beta$ -caryophyllene oxide isolated from Jeju guava (*Psidium cattleianum* Sabine) leaf. *Records of Natural Products*, 2011; 5: 242-246.
21. Legault J, Dahl W, Debiton E, Pichette A, Madelmont JC. Antitumor activity of balsam fir oil: Production of reactive oxygen species induced by  $\alpha$ -humulene as possible mechanism of action. *Planta Medica*, 2003; 69: 402-407.
  22. Sylvestre M, Pichette A, Lavoie S, Longtin A, Legault J. Composition and cytotoxic activity of the leaf essential oil of *Comptonia peregrina* (L.) Coulter. *Phytotherapy Research*, 2007; 21: 536-540.
  23. Duh CY, Wang SK, Weng YL, Chiang MY, Dai CF. Cytotoxic terpenoids from the Formosan soft coral *Nephthea brassica*. *Journal of Natural Products*, 1999; 62: 1518-1521.
  24. Li X, Wang G, Zhao J, Ding H, Cunningham C, Chen F, Flynn DC, Reed E, Li QQ. Antiproliferative effect of  $\beta$ -elemene in chemoresistant ovarian carcinoma cells is mediated through arrest of the cell cycle at the G2-M phase. *Cellular and Molecular Life Sciences*, 2005; 62: 894-904.
  25. Tao L, Zhou L, Zheng L, Yao M. Elemene displays anti-cancer ability on laryngeal cancer cells in vitro and in vivo. *Cancer Chemotherapy and Pharmacology*, 2006; 58: 24-34.