



ISSN: 2321 9114
 AJEONP 2015; 2(5): 34-37
 © 2015 AkiNik Publications
 Received: 15-05-2015
 Accepted: 17-06-2015

Noura S. Dosoky
 Department of Chemistry,
 University of Alabama in Huntsville
 Huntsville, AL 35899, USA

Suraj K. Pokharel
 Department of Chemistry,
 University of Alabama in Huntsville
 Huntsville, AL 35899, USA

William N. Setzer
 Department of Chemistry,
 University of Alabama in Huntsville
 Huntsville, AL 35899, USA

Leaf essential oil composition, antimicrobial and cytotoxic activities of *Cleistocalyx operculatus* from Hetauda, Nepal

Noura S. Dosoky, Suraj K. Pokharel, William N. Setzer

Abstract

The present study investigated the chemical composition and bioactivity of the leaf essential oil from *Cleistocalyx operculatus* (Roxb.) Merr. & L.M. Perry from Nepal. In this work, leaf essential oil of *C. operculatus* was obtained by hydrodistillation and analyzed by GC-MS. The major component of the leaf oil was myrcene (69.7%), followed by (*E*)- β -ocimene (12.24%), (*Z*)- β -ocimene (4.79%), and linalool (4.08%), with other minor constituents. Antimicrobial activities were determined using the microbroth dilution technique, and *in-vitro* cytotoxic activity against MCF-7, MDA-MB-231 and 5637 cells was tested using the MTT assay. *C. operculatus* oil has shown no antimicrobial or cytotoxic activity.

Keywords: *Cleistocalyx operculatus*, GC-MS, essential oil composition, myrcene.

1. Introduction

Cleistocalyx operculatus (Roxb.) Merr. & L.M. Perry (Myrtaceae) is a well-known evergreen perennial tree. It is also known as *Cleistocalyx cerasoides* (Roxb.) I.M. Turner, *Syzygium nervosum* DC, *S. operculatum* (Roxb.) Nied, and *Eugenia operculata* Roxb^[1]. It is widely distributed in India, Nepal, Indonesia, Malaysia, Myanmar, Sri Lanka, Thailand, Vietnam, and Australia. In Nepal, it is commonly known as “kyaman”. The common English name is “sugar root”^[2]. The average tree can reach up to 15 m high and 8-12 m wide. The trunk is smooth, covered with thin brown or greyish brown, flaky bark^[1]. Leaves are simple, opposite, elliptical to obovate, leathery, coarsely veined, measuring 8-10 cm long and 4.5-7 cm wide^[3]. When crushed, leaves emit a mango-like odor. Flowers arise on lateral branches, in 6-12 cm long panicles, collecting the flowers in sessile terminal heads as a compound cyme. Pale yellow flowers are enclosed in the calyx^[1, 4, 5]. Each inflorescence has an average of 30-40 flowers. There are numerous stamens and one pistil. Fruit is a fleshy berry, ovoid, purple to black when ripe^[6]. *C. operculatus* blossoms in March and April and the seeds ripen in May and June. For many years, leaves and bark of *C. operculatus* have been used as a tea to treat stomachache and gastrointestinal disorders as well as an antiseptic for dermatophytic disorders^[6]. The bark decoction is used in taking bath against sunstroke and to wash wounds. A paste of bark and leaves mixed with *Glycyrrhiza glabra* is used to treat pneumonia^[2]. Previous reports showed that the bark of *C. operculatus* contains oleanane-type triterpenes with antidermatophytic activity^[7], and the buds contain flavonoids and triterpene acids^[8]. The essential oil from the buds of *C. operculatus* from Vietnam has previously shown antimicrobial activity against several bacterial strains, including methicillin-resistant *Staphylococcus aureus* (MRSA)^[9] and *Xanthomonas campestris*^[10]. The components of leaf essential oil of *C. operculatus* from Vietnam have been reported^[11]. However, the constituents of plant essential oils can differ from one place to another. The chemical composition and yield of essential oils are influenced by provenance, weather, soil conditions, time of harvest and the drying technique^[12]. Over time, these factors can encourage the biosynthesis of certain terpenes and stop the synthesis of others which subsequently can change the chemical profile of plant essential oils^[13, 14]. To the best of our knowledge, the chemical composition of leaf essential oil of *C. operculatus* from Hetauda, Nepal has not been studied yet, nor have its antimicrobial and cytotoxic activities been investigated.

Correspondence
William N. Setzer
 Department of Chemistry,
 University of Alabama in Huntsville
 Huntsville, AL 35899, USA

2. Materials and Methods

2.1 Plant Material

The leaves of *Cleistocalyx operculatus* were collected from Makwanpur district, Hetauda, Nepal (27°25'N 85°02'E, 550 m asl) in August 2014. The plant was collected and identified by Kiran K. Pokharel, and a voucher specimen (M2013.01) has been deposited with the Department of Forest Research and Survey, Ministry of Forest Research and Soil Conservation, Babarmahal, Kathmandu. The plant material was air-dried for several days. A sample of 100 g was hydrodistilled using a Likens-Nickerson apparatus with continuous extraction using chloroform for 4 h to give 0.4 mL of yellow essential oil with pleasant odor, which was stored at 4 °C until analysis.

2.2 Gas Chromatographic – Mass Spectral Analysis

The essential oil of *C. operculatus* was analyzed by GC-MS using an Agilent 6890 GC with Agilent 5973 mass selective detector, an HP-5ms fused silica capillary column and an Agilent Chem Station data system as previously described [15]. Identification of the oil components was based on their retention indices (RI) and by comparison of their mass spectral fragmentation patterns with those reported in the literature [16].

2.3 Antimicrobial Screening

The essential oil of *C. operculatus* was screened for antimicrobial activity against *Bacillus cereus* (ATCC No. 14579), *Staphylococcus aureus* (ATCC No. 29213), *Aspergillus niger* (ATCC No. 16888) and *Candida albicans* (ATCC No.10231). The minimum inhibitory concentration (MIC) was determined using the microbroth dilution technique as previously reported [17].

2.4 Cytotoxicity Screening

MCF-7 human breast adenocarcinoma cells (ATCC No. HTB-22) and MDA-MB-231 cells (ATCC No. HTB-26) were grown in RPMI 1640 supplemented with 10% Fetal bovine serum (FBS), 30 mM HEPES, NaHCO₃, and Penicillin-Streptomycin. 5637 human bladder cancer cells (ATCC No. HTB-9) were grown in RPMI 1640 supplemented with 10% FBS, 1 mM sodium pyruvate, 2.5 g/L glucose, 30 mM HEPES, NaHCO₃, and Penicillin-Streptomycin. *In-vitro* cytotoxic activity of *C. operculatus* oil on MCF-7, MDA-MB231 and 5637 cells was performed using the 96-well MTT assay as previously reported [18].

3. Results and Discussion

Monoterpenes predominated in *C. operculatus* essential oil composition as shown in Table 1. The essential oil is mainly composed of myrcene (69.7%), followed by (*E*)- β -ocimene (12.24%), (*Z*)- β -ocimene (4.79%), and linalool (4.08%), with other minor constituents (3% or less of each). The essential oil composition and percentages in the current study are significantly different from the previously published report from Vietnam. The major components of leaf essential oil of *C. operculatus* of Vietnamese origin were reported to be (*Z*)- β -ocimene (32.1%), myrcene (24.6%), and (*E*)-caryophyllene (14.5%) and (*E*)- β -ocimene (9.4%) [11]. This difference might be due to several factors including age, vegetative cycle stage, climate, season, soil composition and edaphic factors [19]. Also, the essential oil composition of *C. operculatus* is different from the leaf essential oil of *Syzygium aromaticum* and *S. cumini*. *S. aromaticum* had eugenol (94.4%) as the main constituent followed by (*E*)-caryophyllene (2.9%) [20] While the main oil constitutes in *S. cumini* were α -pinene (17.53%), α -terpineol (16.67 %) and *allo*-ocimene (13.55%) [21].

Table 1: Chemical composition of *Cleistocalyx operculatus* leaf oil from Nepal.

RI ^a	Compound	% \pm SD ^b
934	α -Thujene	0.53 \pm 0.03
992	Myrcene	69.70 \pm 1.02
1003	α -Phellandrene	0.69 \pm 0.02
1023	<i>p</i> -Cymene	2.71 \pm 0.06
1027	β -Phellandrene	1.20 \pm 0.05
1037	(<i>Z</i>)- β -Ocimene	4.79 \pm 0.06
1047	(<i>E</i>)- β -Ocimene	12.24 \pm 0.07
1057	γ -Terpinene	0.82 \pm 0.04
1086	Terpinolene	0.26 \pm 0.03
1098	Linalool	4.08 \pm 0.20
1174	Terpinen-4-ol	tr ^c
1416	(<i>E</i>)-Caryophyllene	1.11 \pm 0.17
1562	(<i>E</i>)-Nerolidol	1.57 \pm 0.27
1521	Caryophyllene oxide	0.23 \pm 0.09
1620	(<i>Z</i>)-Asarone	0.05 \pm 0.08
	Compounds Identified	15 (99.72%)

^a RI determined with respect to a homologous series of *n*-alkanes on an HP-5ms column.

^b Average of three measurements \pm standard deviations.

^c tr = "trace" (< 0.05%).

The most abundant constituent was myrcene (7-methyl-3-methylene-1, 6-octadiene), a well-known monoterpene. It was also reported in the essential oils of several plant species such as *Myrcia* spp. [22], *Porophyllum ruderale* [23], fresh mango fruit [24], *Cannabis sativa* [25], *Pimenta racemosa* [26], *Citrus aurantium* [27], *Humulus lupulus* [28], *Verbena* spp., and *Cymbopogon citratus* [29]. Due to its muscle-relaxing, anti-depressant, anti-inflammatory [23], anti-tumor and analgesic effects [30], myrcene is used in the treatment of spasms, insomnia, and pain. It was also used to minimize epileptic seizures [31] and to prevent peptic ulcer disease [32]. In addition, myrcene was found to affect the integrity and permeability of the cell membrane [33]. Myrcene is also very important in perfumery because of its pleasant odor, which is described as earthy, clove-like, and fruity. It is used as an intermediate in the commercial production of various compounds like geraniol, nerol, linalool, myrcenol, citral, citronellal, hydroxycitronellal, menthol, and ionones [34]. Based on our results, the mature leaves of *C. operculatus* from Nepal can be considered as a good source for commercial isolation of natural myrcene.

The essential oil of *C. operculatus* was screened for potential antimicrobial activity (Table 2). The results showed that there was no antibacterial activity against *Bacillus cereus* (MIC = 1250 μ g/mL) or *Staphylococcus aureus* (MIC = 625 μ g/mL) and no antifungal activity against *Aspergillus niger* (MIC = 625 μ g/mL) and *Candida albicans* (MIC = 2500 μ g/mL). The essential oil was screened for *in-vitro* cytotoxic activity against MCF-7, MDA-MB-231 and 5637 cells but showed no significant activity (0% kill at 100 μ g/mL).

Table 2: Antimicrobial activity (MIC, μ g/mL) of *Cleistocalyx operculatus* leaf oil from Nepal.

<i>B. cereus</i>	<i>S. aureus</i>	<i>A. niger</i>	<i>C. albicans</i>
1250	625	625	2500

4. Conclusions

The essential oil of *Cleistocalyx operculatus* from Nepal was analyzed by GC-MS and was found to be a good source of myrcene (69.7%) followed by (*E*)- β -ocimene (12.24%), (*Z*)- β -ocimene (4.79%), and linalool (4.08%). *C. operculatus*

essential oil was screened for antimicrobial activity and cytotoxic activity but was found to be inactive.

5. Acknowledgments

We are grateful to Kiran K. Pokharel (Assistant Research Officer in the Department of Forest Research and Survey, Ministry of Forest Research and Soil Conservation, Babarmahal, Kathmandu, Nepal) for identification and collection of plant materials.

6. References

- Zhengyi W, Raven PH, Deyuan H. Flora of China, Clusiaceae through Araliaceae. Missouri Botanical Garden Press 2008; 13:344.
- Quattrocchi U. CRC World Dictionary of Medicinal and Poisonous Plants: Common Names, Scientific Names, Eponyms, Synonyms, and Etymology. CRC Press, 2012, 3641-3642.
- Nguyen, Duong Van. Medicinal Plants of Vietnam, Cambodia, and Laos. Santa Monica, CA: Mekong Printing, 1993.
- Roxburgh W. *Flora Indica*, or, Descriptions of Indian Plants. Volume II. Serampore: Printed for W. Thacker and Co, Calcutta and Parbury, Allen and Co, London, 1832, 486.
- Thongma S. Botanical description of Makiang. Makiang. Lampang Agricultural Research and Training Center, Bangkok, 2002, 66-96.
- Loi DT. Vietnamese Medicinal Plants and Herbs. Hanoi Pub. House Techno. Vietnam, 1986, 437-438.
- Nomura M, Yamakawa K, Hirata Y, Niwa M. Antidermatophytic constituent from the bark of *Cleistocalyx operculatus*. Shoyakugaku Zasshi 1993; 47:408-410.
- Zhang FX, Liu MF, Lu RR. Chemical constituents from the buds of *Cleistocalyx operculatus*. Zhiwu Xuebao 1990; 32:469-472.
- Dung NT, Kim JM, Kang SC. Chemical composition, antimicrobial and antioxidant activities of the essential oil and the ethanol extract of *Cleistocalyx operculatus* (Roxb.) Merr and Perry buds. Food and Chemical Toxicology 2008; 46:3632-3639.
- Bajpai VK, Dung NT, Suh HJ, Kang SC. Antibacterial activity of essential oil and extracts of *Cleistocalyx operculatus* buds against the bacteria of *Xanthomonas* spp. Journal of the American Oil Chemists' Society. 2010; 87:1341-1349.
- Dung NX, Luu HV, Khoi TT. GC and GC/MS analysis of the leaf oil of *Cleistocalyx operculatus* Roxb. Merr. et Perry (Syn. *Eugenia operculata* Roxb.; *Syzygium nervosum* DC). Journal of Essential Oil Research. 1994; 6:661-662.
- Boira H, Blanquer A. Environmental factors affecting chemical variability of essential oils in *Thymus piperella* L. Biochemical Systematics and Ecology 1997; 26:811-822.
- Rajeswara Rao BR, Kaul PN, Mallavarapu GR, Ramesh S. Effect of seasonal climatic changes on biomass yield and terpenoid composition of rosescented geranium. Biochemical Systematics and Ecology 1996; 24:627-635.
- Loziene K, Venskutonis PR. Influence of environmental and genetic factors on the stability of essential oil composition of *Thymus pulegioides*. Biochemical Systematics and Ecology, 2004; 33:517-525.
- Monzote L, Nance MR, Garcia M, Scull R, Setzer WN. Comparative chemical, cytotoxicity and antileishmanial properties of essential oils from *Chenopodium ambrosioides*. Natural Product Communications 2011; 6:281-286.
- Adams RP. Identification of Essential Oil Components by Gas Chromatography/Mass Spectrometry, 4th Ed. Allured Publishing, Carol Stream, Illinois, 2007.
- Setzer MC, Setzer WN, Jackes BR, Gentry GA, Moriarity DM. The medicinal value of tropical rainforest plants from Paluma, North Queensland, Australia. Pharmaceutical Biology 2001; 39:67-78.
- Palazzo MC, Wright HL, Agius BR, Wright BS, Moriarity DM, Haber WA *et al.* Chemical compositions and biological activities of leaf essential oils of six species of Annonaceae from Monteverde, Costa Rica. Records of Natural Products 2009; 3:153-160.
- Angioni A, Barra A, Coroneo V, Dessi S, Cabras P. Chemical composition, seasonal variability, and antifungal activity of *Lavandula stoechas* L. ssp. *stoechas* essential oils from stem/leaves and flowers. Journal of Agricultural and Food Chemistry. 2006; 54:4364-4370.
- Raina VK, Srivastava SK, Aggarwal KK, Syamasundar KV, Kumar S. Essential oil composition of *Syzygium aromaticum* leaf from Little Andaman, India. Flavour and Fragrance Journal. 2001; 16:334-336.
- Elansary HO, Salem MZM, Ashmawy NA, Yacout MM. Chemical composition, antibacterial and antioxidant activities of leaves essential oils from *Syzygium cumini* L., *Cupressus sempervirens* L. and *Lantana camara* L. from Egypt. Journal of Agricultural Science. 2012; 4(10):144-152.
- Zoghbi MdGB, Andrade EHA, da Silva MHL, Carreira LMM, Maia JGS. Essential oils from three *Myrcia* species. Flavour and Fragrance Journal. 2003; 18:421-424.
- Souza MC, Siani AC, Ramos MFS, Menezes-de-Lima Jr O, Henriques MGMO. Evaluation of anti-inflammatory activity of essential oils from two Asteraceae species. Pharmazie, 2003; 58:582-586.
- Pandit SS, Chidley HG, Kulkarni RS, Pujari KH, Giri AP, Gupta VS. Cultivar relationships in mango based on fruit volatile profiles. Food Chemistry, 2009; 114(1):363-372.
- Mediavilla V, Steinemann S. Essential oil of *Cannabis sativa* L. strains. Journal of the International Hemp Association. 1997; 4(2):80-82.
- Attokaran M. Natural Food Flavors and Colorants. John Wiley & Sons, 2011, 14.
- Moraes TM, Kushima H, Moleiro FC, Santos RC, Rocha LRM *et al.* Effects of limonene and essential oil from *Citrus aurantium* on gastric mucosa: role of prostaglandins and gastric mucus secretion, Chemo-Biological Interactions. 2009; 180:499-505.
- Nance MR, Setzer WN. Volatile components of aroma hops (*Humulus lupulus* L.) commonly used in beer brewing. Journal of Brewing and Distilling. 2011; 2(2):16-22.
- Günther E, Althausen D. β -myrcene, in: E. Günther (Ed.), The Essential Oils. The constituents of essential oils. New York, 1949, 8-10.
- Rao VS, Menezes AM, Viana GS. Effect of myrcene on nociception in mice. Journal of Pharmacy and Pharmacology. 1990; 42(12):877-878.

31. Elisabetsky E, de Souza GPC, dos Santos MAC, Siqueira IR, Amador TA, Nunes DS. Sedative properties of linalool. *Fitoterapia*, 1995; 66:407-414.
32. Bonamin F, Moraes TM, dos Santos RC, Kushima H, Faria FM, Silva MA *et al.* The effect of a minor constituent of essential oil from *Citrus aurantium*: The role of β -myrcene in preventing peptic ulcer disease. *Chemico-Biological Interactions*. 2014; 212:11-19.
33. Cox SD, Markham JL. Susceptibility and intrinsic tolerance of *Pseudomonas aeruginosa* to selected plant volatile compounds. *Journal of Applied Microbiology*. 2007; 103:930-936.
34. Leung AY. *Encyclopedia of Common Natural Ingredients Used in Food, Drugs and Cosmetics*, 3rd Ed., Wiley, New York, 1980.