



AkiNik

American Journal of Essential Oils and Natural Products

Available online at www.essencejournal.com

A
J
E
O
N
P

American
Journal of
Essential
Oils and
Natural
Products

ISSN: 2321 9114

AJEONP 2013; 1 (1): 48-50

© 2013 AkiNik Publications

Received 24-8-2013

Accepted: 16-9-2013

1-Nitro-2-phenylethane Dominates the Chemical Composition of the Leaf Essential Oil of *Uvaria chamae* from Badagry, Nigeria

Moses S. Owolabi*, Olowu R. Adewale, Noura S. Dosoky and William N. Setzer

ABSTRACT

The leaf essential oil of *Uvaria chamae* (Annonaceae) was obtained by hydrodistillation and analyzed by GC-MS. A total of 69 compounds were identified in the leaf oil (99.6% of the composition), which was dominated by 1-nitro-2-phenylethane (63.2%), linalool (9.9%), and germacrene D (6.6%). The bioactivities of the major constituents account for the traditional uses of *U. chamae* leaves to treat fevers, wounds, swellings, and injuries.

Keywords: *Uvaria chamae*, Essential Oil Composition, 1-nitro-2-phenylethane, Linalool

1. Introduction

The Annonaceae is a large family of shrubby aromatic plants composed of 112 genera with about 2,150 species [1]. *Uvaria chamae* P. Beauv., commonly known as “finger root” or “banana”, is a climbing shrub that grows to about 3.6 m to 4.5 m tall and is found in the tropical wet and dry forests of west and central Africa along coastal scrubland. The fruit is edible and widely eaten, and the plant has been used for treatment of fevers, tumor growth, stroke and cases of venereal disease [2, 3]. The leaves of *U. chamae* have been used to treat wounds and sores, injuries, swellings, and to treat “yellow fever” [4, 5]. Leaf extracts of *U. chamae* have been reported to possess antibiotic, antispasmodic and anti-inflammatory properties [3]. Previous studies of *U. chamae* have shown the plant to be rich in sesquiterpene hydrocarbons dominated by germacrene D and γ -cadinene [6, 7]. In this report we present the chemical characterization of the leaf essential oil of *U. chamae* collected from Badary, Nigeria.

2. Materials and Methods

2.1 Plant Material

Leaves of *U. chamae* were collected from a mature tree on April, 2011. The plant was identified at the University of Lagos herbarium and a voucher specimen (LUH 3626) has been deposited in the herbarium of the Botany Department. The leaves of the plant (350 g) were air-dried, chopped and hydrodistilled using Clevenger apparatus [8] to yield 0.75 g light yellow essential oil. The essential oil so obtained was stored in a sealed glass bottle with screw lid cover under refrigeration at 4 °C.

2.2 GC-MS Analysis

The volatile oil sample was subjected to GC-MS analysis on an Agilent system consisting of an Agilent model 6890 Gas Chromatograph, an Agilent 5973 mass selective detector (EIMS, electron energy = 70 eV, scan range = 45-400 amu, and scan rate = 3.99 scans/sec) and an Agilent Chemstation data system. The GC column was a HP-5ms fused silica capillary with a (5% phenyl)-methylpolysiloxane stationary phase, film thickness 0.25 μ m, length 30 m, and internal diameter of 0.25 mm. The carrier gas was helium with a column head pressure of 7.07 psi and a flow rate of 1.0 mL/min. Inlet temperature was 200 °C and MSD detector temperature was 280 °C. The GC oven temperature program was used as follows: 40 °C initial temperature, hold for 10 min, increased at 3 °C/min to 200 °C, increased 2 °C/min to 220 °C.

Correspondence:

Moses S. Owolabi

Department of Chemistry, Lagos State University, P.M.B 001, Ojo, Lagos, Nigeria.

E-mail:sunnyconcept2007@yahoo.com

The sample was dissolved in dichloromethane and a splitless injection technique was used. Identification of the constituents of the volatile oil was achieved based on their retention data (retention indices) determined with reference to C₉-C₂₁ *n*-alkane homologous series, and by comparison of their mass spectral fragmentation patterns with those reported in the literature¹⁹ and stored on the

MS library [NIST database (G1036A, revision D.01.00) / ChemStation data system (G1701CA, version C.00.01.08)]. The chemical composition of *Uvaria chamae* essential oil is summarized in Table 1.

Table 1: Chemical Composition of *Uvaria chamae* Leaf Essential Oil

RI	Compound	%	RI	Compound	%
941	α-Pinene	0.3	1459	(<i>E</i>)-β-Farnesene	0.2
953	Camphene	Tr	1460	Alloaromadendrene	0.1
964	Benzaldehyde	Tr	1463	<i>cis</i> -Cadina-1(6),4-diene	tr
978	β-Pinene	0.2	1474	<i>trans</i> -Cadina-1(6),4-diene	0.1
992	Myrcene	0.2	1479	γ-Murolene	0.2
1004	α-Phellandrene	Tr	1483	Germacrene D	6.6
1010	δ-3-Carene	Tr	1487	β-Selinene	1.0
1016	α-Terpinene	Tr	1492	<i>trans</i> -Muurola-4(14),5-diene	0.3
1024	<i>p</i> -Cymene	0.4	1495	<i>epi</i> -Cubebol	0.7
1028	Limonene	0.1	1497	Bicyclogermacrene	0.3
1030	1,8-Cineole	0.6	1501	α-Murolene	0.3
1038	(<i>Z</i>)- <i>b</i> -Ocimene	0.1	1505	Germacrene A	0.2
1049	(<i>E</i>)- <i>b</i> -Ocimene	1.9	1511	(<i>E,E</i>)-α-Farnesene	0.1
1058	γ-Terpinene	0.2	1516	Cubebol	0.8
1100	Linalool	9.9	1525	δ-Cadinene	2.1
1139	Benzeneacetonitrile	0.1	1533	<i>trans</i> -Cadina-1,4-diene	0.1
1143	(<i>E</i>)-Epoxyocimene	Tr	1537	α-Cadinene	tr
1164	Borneol	0.1	1550	Elemol	0.1
1176	Terpinen-4-ol	Tr	1557	Germacrene B	tr
1189	α-Terpineol	0.1	1566	(<i>E</i>)-Nerolidol	1.4
1231	(3 <i>Z</i>)-Hexenyl 2-methylbutanoate	Tr	1570	(3 <i>Z</i>)-Hexenyl benzoate	0.2
1285	Bornyl acetate	Tr	1578	Spathulenol	0.2
1296	1-Nitro-2-phenylethane	63.2	1583	Caryophyllene oxide	0.3
1325	(3 <i>Z</i>)-Hexenyltiglate	0.1	1582	Gleenol	tr
1349	α-Cubebene	0.2	1610	Humulene epoxide II	0.1
1375	α-Copaene	0.4	1628	1- <i>epi</i> -Cubenol	0.4
1384	β-Bourbonene	0.2	1641	τ-Cadinol	tr
1390	β-Cubebene	0.2	1643	τ-Murolol	0.7
1392	β-Elemene	0.2	1646	α-Murolol (= Torreyol)	0.2
1419	(<i>E</i>)-Caryophyllene	1.7	1655	α-Cadinol	0.5
1428	β-Copaene	0.1	1715	Pentadecanal	0.1
1436	α- <i>trans</i> -Bergamotene	0.1	1762	Benzyl benzoate	0.9
1443	6,9-Guaiadiene	Tr	1956	Palmitic acid	0.2
1451	<i>cis</i> -Muurola-3,5-diene	0.1	2108	(<i>E</i>)-Phytol	0.1
1453	α-Humulene	0.6		Total Identified	99.6

3. Results and Discussion

The leaf oil of *U. chamae* in this study was dominated by 1-nitro-2-phenylethane (63.2%) with lesser quantities of linalool (9.9%) and germacrene D (6.6%), and therefore, is notably different compared to previously reported samples [6, 7]. Although nitro-containing compound are relatively rare phytochemicals, 1-nitro-2-phenylethane has been found to be an abundant constituent of *Aniba canelilla* and *Ocotea pretiosa* wood and bark oils [10-12]. 1-Nitro-2-phenylethane has shown anti-inflammatory and antinociceptive activity [13, 14], as well as anticandidal [11] and brine shrimp lethality [15] effects. In addition, linalool has shown anti-inflammatory [16], antinociceptive [17], and antimicrobial [18] activities.

4. Conclusions

The bioactivities of the major constituents of *Uvaria chamae* leaf oil are consistent with the traditional medicinal uses of this plant

5. Reference:

1. Mabberly DJ. The Plant – Book, 2nd Ed. Cambridge University Press, UK, 1997, 41-42.
2. Iwu MM. Handbook of African Medicinal Plants. CRC Press, Boca Raton, Florida, 1993, 145.
3. Okwu DE, Iroabuchi F. Phytochemical composition and biological activities of *Uvaria chamae* and *Clerodendron splendens*. E.-J. Chem., 2009; 6: 553-560.
4. Ayensu ES. Medicinal Plants of West Africa. Reference Publications, Inc., Algonac, Michigan, 1978, 41.
5. Oliver-Bever B. Medicinal Plants in Tropical West Africa. Cambridge University Press, UK, 1986, 134-135.
6. Ayedoun MA, Moudachirou M, Adeoti BS, Menut C, Lamaty G, Bessière JM. Aromatic plants of tropical West Africa. VII. Essential oil of leaf and root bark of *Uvaria chamae* P. Beauv. from Benin. J. Essent. Oil Res., 1999; 11: 23-26.
7. Oguntimein B, Ekundayo O, Laakso I, Hiltunen I. Volatile constituents of *Uvaria chamae* leaves and root bark. Planta Med., 1989; 55: 312-313.
8. British Pharmacopoeia. Vol. II. H.M. Stationary office, London, 1980, A 109.
9. Adams R P. Identification of Essential Oil Components by Gas Chromatography / Mass Spectrometry, 4th Ed. Allured Publishing Corp, Carol Stream, Illinois, 2007.
10. Gottlieb OR, Magalhães MT. Occurrence of 1-nitro-2-phenylethane in *Ocotea pretiosa* and *Aniba canelilla*. J. Org. Chem., 1959; 24: 2070-2071.
11. Oger JM, Richomme P, Guinaudeau H, Bouchara JP, Fournet A. *Aniba canelilla* (H.B.K.) Mez essential oil: Analysis of chemical constituents, fungistatic properties. J. Essent. Oil Res., 1994; 6: 493-497.
12. Taveira FSN, de Lima WN, Andrade EHA, Maia JGS. Seasonal essential oil variation of *Aniba canelilla*. Biochem. System. Ecol., 2003; 31: 69-75.
13. De Lima AB. Estudo da AçãoAntinociceptiva e Antiinflamatória do 1-Nitro-2-feniletano, PríncipalConstituinte da *Aniba canelilla*. M. S. Thesis, Universidade de Federal do Pará, Belém, Brazil, 2008.
14. de Lima AB, Santana MB, Cardoso AS, da Silva JKR, Maia JGS, Carvalho JCT, Sousa PJC. Antinociceptive activity of 1-nitro-2-phenylethane, the main component of *Aniba canelilla* essential oil. Phytomedicine, 2009; 16: 555-559.
15. Da Silva JKR, Sousa PJC, Andrade EHA, Maia JGS. Antioxidant capacity and cytotoxicity of essential oil and methanol extract of *Aniba canelilla* (H.B.K.) Mez. J. Agric. Food Chem., 2007; 55: 9422-9462.
16. Peana AT, D'Aquila PS, Panin F, Serra G, Pippia P, Moretti MDL. Anti-inflammatory activity of linalool and linalyl acetate constituents of essential oils. Phytomedicine, 2002; 9: 721-726.
17. Peana AT, D'Aquila PS, Chessa ML, Moretti MDL, Serra G, Pippia P. (–)-Linalool produces antinociception in two experimental models of pain. Eur. J. Pharmacol., 2003; 460: 37-41.